



From biology to management: Practical tools for black stem borer control

Dr. Monique J. Rivera and Sandra Lizarraga

Outline

- **Ambrosia beetles in apple overview**
- **Progress on development of a model to predict flight**
- **2025 Results of Verbenone**

Ambrosia beetles in Apple

Apple Tree Damage by Ambrosia Beetles



What ambrosia beetles do we see in apple?



ISSUE: [VOL. 5506 NO. 2: 6 SEPT. 2024](#)

TYPE: ARTICLE

PUBLISHED: 2024-09-06

DOI: [10.11646/ZOOTAXA.5506.2.6](#)

PAGE RANGE: 261-271

ABSTRACT VIEWS: 197





PDF DOWNLOADED: 33

FIRST REPORT OF *ANISANDRUS MAICHE* (COLEOPTERA: CURCULIONIDAE: SCOLYTINAE) INFESTING APPLE TREES


KELSEY N. TOBIN⁺[ID](#), M. ELIZABETH MOORE⁺[ID](#), SANDRA LIZARRAGA⁺[ID](#), JANE PETZOLDT⁺[ID](#), COREY REESE⁺[ID](#), BRIAN LOVETT⁺[ID](#), MONIQUE J. RIVERA⁺[ID](#)

Flight of Foundress Female

1 Beetle = Full Colony within Tree



Ambrosiella grosmanniae



Why are ambrosia beetles attacking apples?

Black stem borer





Photo by Betsy Anderson 2016, *Journal of Integrated Pest Management*



Tree stress → Ethanol and other cues → Ambrosia Beetle Attacks




ABIOTIC

BIOTIC

Younger, smaller trees = More stress

Possible other stress sources: pruning, viral load, topworking

Black Stem Borer Seasonal Flight Patterns



- Two peak flights
- Overwinter as adults
- Can emerge as soon as the weather is suitable



Anisandrus maiche



Zootaxa 5506 (2): 261–271

<https://www.mapress.com/zt/>

Copyright © 2024 Magnolia Press

Article

ISSN 1175-5326 (print edition)

ZOOTAXA

ISSN 1175-5334 (online edition)

<https://doi.org/10.11646/zootaxa.5506.2.6>


<http://zoobank.org/urn:lsid:zoobank.org:pub:497E4655-8462-4B2C-ADA3-1382EE9A8DD8>

FIRST REPORT OF *ANISANDRUS MAICHE* (COLEOPTERA: CURCULIONIDAE: SCOLYTINAE) INFESTING APPLE TREES




KELSEY N. TOBIN^{1*}, M. ELIZABETH MOORE^{2*}, SANDRA LIZARRAGA¹, JANE PETZOLDT¹, COREY REESE², BRIAN LOVETT², MONIQUE J. RIVERA¹




Anisandrus maiche




Ambrosiella cleistominuta







Can we predict flights to optimize application timings?



**Progress on development of a model
to predict Black Stem Borer flight**

Bihourly Trapping Study by Sandra Lizarraga



Five clear sticky trap baited with an Ethanol lure



ibutton sensor to track temperature and humidity


Time Groups (EDT)

8-10	10-12	12-2	2-4	4-6	6-8
AM	PM				

Predicting beetle flight based on environmental factors



Relative humidity



Temperature



Time of day




Victor Alves


PhD candidate of
Plant Pathology at
UC Davis

Predictive Model

A Bayesian model that estimates the optimal environmental conditions for beetle flight using bell-shaped response curves



Results



Optimal relative humidity was estimated at **70.45%**



Activity peaked at a temperature of **75.5F**




Peak time of the day was determined to be at **7:00 PM**

Results

Scaled Predicted Response (0 to 100)

RH fixed at 70 %



- Establishes the first temperature- and time-based flight prediction framework for BSB
- Enables targeted monitoring by improving prediction of first peak flight periods
- Optimizes timing of preventative management actions

Key findings

- Developed a predictive flight scale for *X. germanus*

Implications for apple growers

- Improves decision-making through localized monitoring
- Precision insecticide applications timed with risk

Future Directions

- Expand and refine the model with additional field data
- Get model onto NEWA or some other applet for use




2025 Results of Verbenone Trial

Apple orchard



**Repellent
=
Push**

VERBENONE



Verbenone as a repellent

- Produced in **pine trees** as a response to stress and/or insect attack
- Used as a **repellent** against multiple beetles including ambrosia beetles



Overlap in response to semiochemicals across key pest species



© P. Ziegler



Attraction = 
Repellent = 

Xylosandrus germanus (BSB)

Ethanol 

Verbenone 

Anisandrus maiche

Ethanol 

Verbenone 

Xylosandrus crassiusculus

Ethanol 

Verbenone 

Comparing commercially available verbenone products



◆ Verbenone sachets

- 100 g of verbenone
- 15 sachets (\$147 USD)
- ¼-acre plots
- 8 weeks of coverage



⦿ SPLAT Verbenone


- 700 g of verbenone
- 1 canister (\$122 USD)
- 1-acre plot
- 12-16 weeks of coverage
- State-by-state regulations
- Food-safe, biodegradable



● Test SPLAT

- New SPLAT product
- Currently non-commercial
- Designed for extended field longevity

Comparison of commercially available verbenone dispensers



● Test SPLAT

◆ Verbenone Pouch


▲ Low-release EtOH trap (monitoring)

○ SPLAT VERB



Species	New York
<i>X. germanus</i>	60
<i>A. maiche</i>	95
<i>X. crassiusculus</i>	1


Mean Beetle Captures - New York




All verbenone formulations were successful in reducing ambrosia beetle populations



Mean Beetle Captures - Pennsylvania



Species	Pennsylvania
<i>X. germanus</i>	24
<i>A. maiche</i>	91
<i>X. crassiusculus</i>	838



All verbenone formulations were successful in reducing ambrosia beetle populations even under high beetle pressure

Key Finding

- All verbenone dispensers reduce ambrosia beetle capture in treated blocks

Future Directions

- Assess longevity of verbenone and phytotoxicity





Thank You

Questions?